


19. Let the curve $z(1+i) + \bar{z}(1-i) = 4$, $z \in \mathbb{C}$, divide the region $|z-3| \leq 1$ into two parts of areas α and β . Then $|\alpha - \beta|$ equals :

(1) $1 + \frac{\pi}{2}$ (2) $1 + \frac{\pi}{3}$
 (3) $1 + \frac{\pi}{4}$ (4) $1 + \frac{\pi}{6}$

Ans. (1)



Let $z = x + iy$

$$(x+iy)(1+i) + (x-iy)(1-i) = 4$$

$$x+ix+iy-y+x-ix-iy-y=4$$

$$2x-2y=4$$

$$x-y=2$$

$$|z-3| \leq 1$$

$$(x-3)^2 + y^2 \leq 1$$

$$\text{Area of shaded region} = \frac{\pi \cdot 1^2}{4} - \frac{1}{2} \cdot 1 \cdot 1 = \frac{\pi}{4} - \frac{1}{2}$$

Area of unshaded region inside the circle

$$= \frac{3}{4} \pi \cdot 1^2 + \frac{1}{2} \cdot 1 \cdot 1 = \frac{3\pi}{4} + \frac{1}{2}$$

$$\therefore \text{difference of area} = \left(\frac{3\pi}{4} + \frac{1}{2} \right) - \left(\frac{\pi}{4} - \frac{1}{2} \right)$$


$$= \frac{\pi}{2} + 1$$

20. The area of the region enclosed by the curves $y = x^2 - 4x + 4$ and $y^2 = 16 - 8x$ is :

(1) $\frac{8}{3}$ (2) $\frac{4}{3}$
 (3) 5 (4) 8

Ans. (1)

Sol.



$$y = (x-2)^2, y^2 = 8(x-2)$$

$$y = x^2, y^2 = -8x$$

$$= \frac{16ab}{3} = \frac{16 \times \frac{1}{4} \times 2}{3} = \frac{8}{3}$$

SECTION-B

21. Let $y = f(x)$ be the solution of the differential

$$\text{equation } \frac{dy}{dx} + \frac{xy}{x^2-1} = \frac{x^6+4x}{\sqrt{1-x^2}}, -1 < x < 1 \text{ such}$$

that $f(0) = 0$. If $6 \int_{-\frac{1}{2}}^{\frac{1}{2}} f(x) dx = 2\pi - \alpha$ then α^2 is

equal to _____.

Ans. (27)

Sol. I.F. $e^{-\frac{1}{2} \int \frac{2x}{1-x^2} dx} = e^{-\frac{1}{2} \ln(1-x^2)} = \sqrt{1-x^2}$

$$y \times \sqrt{1-x^2} = \int (x^6 + 4x) dx = \frac{x^7}{7} + 2x^2 + c$$

$$\text{Given } y(0) = 0 \Rightarrow c = 0$$

$$y = \frac{\frac{x^7}{7} + 2x^2}{\sqrt{1-x^2}}$$

$$\text{Now, } 6 \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\frac{x^7}{7} + 2x^2}{\sqrt{1-x^2}} dx = 6 \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{2x^2}{\sqrt{1-x^2}} dx$$

$$= 24 \int_0^{\frac{1}{2}} \frac{x^2}{\sqrt{1-x^2}} dx$$

Put $x = \sin\theta$

$$dx = \cos\theta d\theta$$

$$= 24 \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta$$

$$= 24 \int_0^{\frac{\pi}{6}} \left(\frac{1-\cos 2\theta}{2} \right) d\theta = 12 \left[\theta - \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{6}}$$

$$= 12 \left(\frac{\pi}{6} - \frac{\sqrt{3}}{4} \right)$$


$$= 2\pi - 3\sqrt{3}$$

$$\alpha^2 = (3\sqrt{3})^2 = 27$$

22. Let $A(6, 8)$, $B(10 \cos\alpha, -10 \sin\alpha)$ and $C(-10 \sin\alpha, 10 \cos\alpha)$, be the vertices of a triangle. If $L(a, 9)$ and $G(h, k)$ be its orthocenter and centroid respectively, then $(5a - 3h + 100 \sin 2\alpha)$ is equal to _____

Ans. (145)

Sol. All the three points A, B, C lie on the circle $x^2 + y^2 = 100$ so circumcentre is $(0, 0)$



$$\frac{a+0}{3} = h \Rightarrow a = 3h$$

$$\text{and } \frac{9+0}{3} = k \Rightarrow k = 3$$

$$\text{also centroid } \frac{6+10\cos\alpha-10\sin\alpha}{3} = h$$

$$\Rightarrow 10(\cos\alpha - \sin\alpha) = 3h - 6 \quad \dots\text{(i)}$$

$$\text{and } \frac{8+10\cos\alpha-10\sin\alpha}{3} = k$$

$$\Rightarrow 10(\cos\alpha - \sin\alpha) = 3k - 8 = 9 - 8 = 1 \dots\text{(ii)}$$

on squaring $100(1 - \sin 2\alpha) = 1$

$$\Rightarrow 100\sin 2\alpha = 99$$

$$\text{from equ. (i) and (ii) we get } h = \frac{7}{3}$$

$$\text{Now } 5a - 3h + 6k + 100\sin 2\alpha$$


$$= 15h - 3h + 6k + 100\sin 2\alpha$$

$$= 12 \times \frac{7}{3} + 18 + 99$$

$$= 145$$

23. Let the distance between two parallel lines be 5 units and a point P lie between the lines at a unit distance from one of them. An equilateral triangle PQR is formed such that Q lies on one of the parallel lines, while R lies on the other. Then $(QR)^2$ is equal to _____.

Ans. (28)



$$PR = \operatorname{cosec} \theta, PQ = 4 \sec(30 + \theta)$$

For equilateral

$$d = PR = PQ$$

$$\Rightarrow \cos(\theta + 30^\circ) = 4 \sin \theta$$

$$\Rightarrow \frac{\sqrt{3}}{2} \cos \theta - \frac{1}{2} \sin \theta = 4 \sin \theta$$

$$\Rightarrow \tan \theta = \frac{1}{3\sqrt{3}}$$

$$QR^2 = d^2 = \operatorname{cosec}^2 \theta = 28$$

24. If $\sum_{r=1}^{30} \frac{r^2 \binom{30}{r}}{\binom{30}{r-1}} = \alpha \times 2^{29}$, then α is equal to _____.

Ans. (465)

$$\begin{aligned} \text{Sol. } & \sum_{r=1}^{30} \frac{r^2 \binom{30}{r}}{\binom{30}{r-1}} \\ &= \sum_{r=1}^{30} r^2 \left(\frac{31-r}{r} \right) \cdot \frac{30!}{r!(30-r)!} \\ & \left(\because \frac{\binom{30}{r}}{\binom{30}{r-1}} = \frac{30-r+1}{r} = \frac{31-r}{r} \right) \\ &= \sum_{r=1}^{30} \frac{(31-r)30!}{(r-1)!(30-r)!} \\ &= 30 \sum_{r=1}^{30} \frac{(31-r)29!}{(r-1)!(30-r)!} \\ &= 30 \sum_{r=1}^{30} (30-r+1)^{29} C_{30-r} \\ &= 30 \left(\sum_{r=1}^{30} (31-r)^{29} C_{30-r} + \sum_{r=1}^{30} 29 C_{30-r} \right) \\ &= 30(29 \times 2^{28} + 2^{29}) = 30(29 + 2)2^{28} \\ &= 15 \times 31 \times 2^{29} \\ &= 465(2^{29}) \\ &\alpha = 465 \end{aligned}$$

25. Let $A = \{1, 2, 3\}$. The number of relations on A, containing (1, 2) and (2, 3), which are reflexive and transitive but not symmetric, is _____.

Ans. (3)

Sol. Transitivity

$$(1, 2) \in R, (2, 3) \in R \Rightarrow (1, 3) \in R$$

For reflexive (1, 1), (2, 2), (3, 3) $\in R$

Now (2, 1), (3, 2), (3, 1)

(3, 1) cannot be taken


(1) (2, 1) taken and (3, 2) not taken

(2) (3, 2) taken and (2, 1) not taken

(3) Both not taken

therefore 3 relations are possible.

30. For a short dipole placed at origin O, the dipole moment P is along x-axis, as shown in the figure. If the electric potential and electric field at A are V_0 and E_0 , respectively, then the correct combination of the electric potential and electric field, respectively, at point B on the y-axis is given by



(1) $\frac{V_0}{2}$ and $\frac{E_0}{16}$ (2) zero and $\frac{E_0}{8}$
 (3) zero and $\frac{E_0}{16}$ (4) V_0 and $\frac{E_0}{4}$

Ans. (3)

Sol. $E_A = \frac{2kP}{r^3} = E_0 \text{ and } V_A = \frac{kP}{r^2} = V_0$

$$E_B = \frac{kP}{(2r)^3} = \frac{E_0}{16} \text{ and } V_B = \frac{k\vec{P} \cdot \hat{r}}{r^2} = 0$$

31. Which one of the following is the correct dimensional formula for the capacitance in F ? M, L, T and C stand for unit of mass, length, time and charge,

(1) $[F] = [C^2 M^{-2} L^2 T^2]$
 (2) $[F] = [C M^{-2} L^{-2} T^{-2}]$
 (3) $[F] = [C M^{-1} L^{-2} T^2]$
 (4) $[F] = [C^2 M^{-1} L^{-2} T^2]$

Ans. (4)

Sol. $C = \frac{q}{V} = \frac{q \cdot q}{V \cdot q} = \frac{q^2}{WD} = \frac{C^2}{ML^2 T^{-2}} = C^2 M^{-1} L^{-2} T^2$

32. An electron projected perpendicular to a uniform magnetic field B moves in a circle. If Bohr's quantization is applicable, then the radius of the electronic orbit in the first excited state is :

(1) $\sqrt{\frac{2h}{\pi eB}}$ (2) $\sqrt{\frac{4h}{\pi eB}}$
 (3) $\sqrt{\frac{h}{2\pi eB}}$ (4) $\sqrt{\frac{h}{\pi eB}}$

Ans. (4)

Sol. $r = \frac{mv}{eB} \text{ and } mvr = \frac{nh}{2\pi} \Rightarrow (eBr)r = \frac{nh}{2\pi}$
 $\Rightarrow r = \sqrt{\frac{nh}{2\pi eB}}$

first excited state : $n = 2 \therefore r = \sqrt{\frac{h}{\pi eB}}$

33. For a diatomic gas, if $\gamma_1 = \left(\frac{C_p}{C_v}\right)$ for rigid molecules and $\gamma_2 = \left(\frac{C_p}{C_v}\right)$ for another diatomic molecules, but also having vibrational modes. Then, which one of the following options is correct ?
 (C_p and C_v are specific heats of the gas at constant pressure and volume)

(1) $\gamma_2 > \gamma_1$ (2) $\gamma_2 = \gamma_1$
 (3) $2\gamma_2 = \gamma_1$ (4) $\gamma_2 < \gamma_1$

Ans. (4)


Sol. $\gamma = \frac{2}{f} + 1$

without vibration : $f = 5 : \gamma_1 = 1.4$

without vibration : $f = 7 : \gamma_2 = 1.14$

$\therefore \gamma_2 < \gamma_1$

34. A rectangular metallic loop is moving out of a uniform magnetic field region to a field free region with a constant speed. When the loop is partially inside the magnetic field, the plot of magnitude of induced emf (ϵ) with time (t) is given by



Ans. (4)

$$B = \text{constant}$$

Motional emf : $\varepsilon = B\ell v = \text{constant}$

Ans. (3)

Sol. $\frac{hc}{\lambda} = \phi + eV \Rightarrow \frac{hc}{\lambda} = 1 + 2 = 3 \text{ eV} \dots\dots(1)$

$$\frac{hc}{\lambda/2} = 6 = 1 + k_{\max} \quad \therefore k_{\max} = 5 \text{ eV}$$

36. Given below are two statements. One is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

Assertion (A) : A simple pendulum is taken to a planet of mass and radius, 4 times and 2 times, respectively, than the Earth. The time period of the pendulum remains same on earth and the planet.

Reason (R) : The mass of the pendulum remains unchanged at Earth and the other planet. In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Both **(A)** and **(R)** are true but **(R)** is NOT the correct explanation of **(A)**
- (2) **(A)** is true but **(R)** is false
- (3) **(A)** is false but **(R)** is true
- (4) Both **(A)** and **(R)** are true and **(R)** is the correct explanation of **(A)**

Ans. (1)

$$\text{Sol. } g = \frac{GM}{R^2}$$


$$g' = \frac{G(4M)}{(2R)^2} = g$$

A is correct, R is correct ; but since $T = 2\pi\sqrt{\frac{\ell}{g}}$

doesn't depend on mass ; R doesn't explain A.

Ans. (3)


$$\text{Sol. } \vec{\tau} = \vec{r} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 2 & 1 & 2 \end{vmatrix} = \hat{i} - 0\hat{j} - \hat{k}$$



A	B	Y
0	0	1
0	1	0
1	0	0
1	1	1

To obtain the given truth table, following logic gate should be placed at G :

NTA Ans. (1)



For NOR gate : $\overline{A\bar{B}} = \bar{A} + B$

	A	B	Y
0	0	0	1
1	0	1	1
1	1	1	1

∴ Bonus

Ans. (2)

$$\text{Sol. } \text{WD} = \vec{F} \cdot \vec{S} = 2 - 2b - 1 = 0$$

$$\therefore b = \frac{1}{2}$$

40. Given below are two statements. One is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

Assertion (A) : In Young's double slit experiment, the fringes produced by red light are closer as compared to those produced by blue light.

Reason (R) : The fringe width is directly proportional to the wavelength of light.

In the light of above statements, choose the **correct** answer from the options given below :


- (1) Both **(A)** and **(R)** are true and **(R)** is the correct explanation of **(A)**
- (2) **(A)** is false but **(R)** is true.
- (3) Both **(A)** and **(R)** are true but **(R)** is **NOT** the correct explanation of **(A)**.
- (4) **(A)** is true but **(R)** is false.

Ans. (2)

$$\text{Sol. } \beta = \frac{\lambda D}{d} \text{ & } \lambda_R > \lambda_b$$

$$\therefore \beta_R > \beta_L$$

Ans. (2)




$$k_i = \frac{1}{2}mv^2$$

$$k_f = \frac{1}{2}m(v \cos 60^\circ)^2 = \frac{1}{8}mv^2$$

$$\Delta k = k_i - k_f = \frac{3}{8}mv^2 = \frac{3}{8} \times 0.1 \times 400 = 15J$$


Ans. (3)



For transmitted green light to be maxima, reflected green should be minima.

$$\Delta P = 2\mu_0 t = n\lambda$$


$$\Rightarrow t = \frac{n\lambda}{2\mu_0} \therefore t_{\min} = \frac{\lambda}{2\mu_0} = \frac{550}{2 \times 2} = 137.5$$



43.

The tube of length L is shown in the figure. The radius of cross section at the point (1) is 2 cm and at the point (2) is 1 cm, respectively. If the velocity of water entering at point (1) is 2 m/s, then velocity of water leaving the point (2) will be :

Ans. (4)



$$A_1 v_1 = A_2 V_2 \Rightarrow 2\pi(2R)^2 = V_2 \pi R^2$$

$$\therefore V_2 = 8 \text{ m/s}$$

44. Given are statements for certain thermodynamic variables,

(A) Internal energy, volume (V) and mass (M) are extensive variables.

(B) Pressure (P), temperature (T) and density (ρ) are intensive variables.

(C) Volume (V), temperature (T) and density (ρ) are intensive variables.

(D) Mass (M), temperature (T) and internal energy are extensive variables.


Choose the **correct** answer from the points given below :

(1) (C) and (D) only (2) (D) and (A) only
(3) (A) and (B) only (4) (B) and (C) only

Ans. (3)

Sol. Extensive variables depends on size or mass of system ex : internal energy, volume, mass

45. A body of mass 100 g is moving in circular path of radius 2 m on vertical plane as shown in figure. The velocity of the body at point A is 10 m/s. The ratio of its kinetic energies at point B and C is :




(Take acceleration due to gravity as 10 m/s^2)

(1) $\frac{2+\sqrt{3}}{3}$ (2) $\frac{2+\sqrt{2}}{3}$
 (3) $\frac{3+\sqrt{3}}{2}$ (4) $\frac{3-\sqrt{2}}{2}$

Ans. (3)

Sol.



$$\frac{1}{2}m \times 100 + 0 = \frac{1}{2}mV_B^2 + mg \left(R - \frac{R\sqrt{3}}{2} \right)$$

$$100 = V_B^2 + 2gR \left(1 - \frac{\sqrt{3}}{2} \right)$$

$$V_B^2 = 100 - 20(2 - \sqrt{3})$$

$$V_B^2 = 60 + 20\sqrt{3})$$

$$K.E_B = \frac{1}{2} m V_B^2 = \frac{m}{2} (60 + 20\sqrt{3})$$

$$\frac{1}{2}m(100) = \frac{1}{2}mV_c^2 + mg\left(\frac{3R}{2}\right)$$

$$100 = V_C^2 = 60$$

$$V_c^2 = 40$$

$$K.E_c = \frac{1}{2} m V_c^2 = \frac{1}{2} m (40)$$

$$K.E_B = \frac{60 + 20\sqrt{3}}{40} = \frac{3}{2} + \frac{\sqrt{3}}{2} = \frac{3 + \sqrt{3}}{2}$$

SECTION-B


46. A proton is moving undeflected in a region of crossed electric and magnetic fields at a constant speed of $2 \times 10^5 \text{ ms}^{-1}$. When the electric field is switched off, the proton moves along a circular path of radius 2 cm. The magnitude of electric field is $x \times 10^4 \text{ N/C}$. the value of x is _____.
Take the mass of the proton = $1.6 \times 10^{-27} \text{ kg}$.

Ans. (2)


Sol. For uniform speed $V = \frac{E}{B}$

$$\begin{aligned} R &= \frac{mV}{eB} \\ &= \frac{mV^2}{eE} \\ \Rightarrow E &= \frac{mV^2}{eR} \\ &= \frac{1.6 \times 10^{-27} \times 4 \times 10^{10}}{1.6 \times 10^{-19} \times 2 \times 10^{-2}} \\ &= 2 \times 10^4 \text{ N/C.} \end{aligned}$$

47. Two long parallel wires X and Y, separated by a distance of 6 cm, carry currents of 5A and 4A, respectively, in opposite directions as shown in the figure. Magnitude of the resultant magnetic field at point P at a distance of 4 cm from wire Y is $x \times 10^{-5} \text{ T}$. The value of x is _____. Take permeability of free space as $\mu_0 = 4\pi \times 10^{-7} \text{ SI units}$.



Ans. (1)



$$\begin{aligned} B &= \frac{\mu_0(5)}{2\pi \times 0.01} - \frac{\mu_0 4}{2\pi \times 0.04} \\ &= -\frac{100\mu_0}{4\pi} \\ &= -100 \times 10^{-7} \\ &= -1 \times 10^{-5} \text{ T} \end{aligned}$$

48. A parallel plate capacitor of area $A = 16 \text{ cm}^2$ and separation between the plates 10 cm, is charged by a DC current. Consider a hypothetical plane surface of area $A_0 = 3.2 \text{ cm}^2$ inside the capacitor and parallel to the plates. At an instant, the current through the circuit is 6A. At the same instant the displacement current through A_0 is _____ mA.

Ans. (1200)

Sol. $J_d = \frac{I}{A} = \frac{6}{16}$

$$\therefore I \text{ through small area} = J_d \times A' = \frac{6}{16} \times 3.2 = 1.2 \text{ A} = 1200 \text{ mA}$$

49. A tube of length 1m is filled completely with an ideal liquid of mass $2M$, and closed at both ends. The tube is rotated uniformly in horizontal plane about one of its ends. If the force exerted by the liquid at the other end is F then angular velocity of the tube is $\sqrt{\frac{F}{\alpha M}}$ in SI unit. The value of α is _____.

Ans. (1)


Sol.




$$F = 2M\omega^2 \frac{\ell}{2} = M\omega^2 \ell$$

$$\omega = \sqrt{\frac{F}{M\ell}}$$

50. The net current flowing in the given circuit is _____ A.



Ans. (1)



Sol.

$$R_{eq} = 2\Omega$$

$$I = \frac{2}{2} = 1 \text{ A}$$

CHEMISTRY

SECTION-A

51. Arrange the following compounds in increasing order of their dipole moment :



(1) NF₃ < HBr < H₂S < CHCl₃

(2) HBr < H₂S < NF₃ < CHCl₃

(3) H₂S < HBr < NF₃ < CHCl₃

(4) CHCl₃ < NF₃ < HBr < H₂S

Ans. (1)




Sol. Increasing order of Dipole moment



$\mu = 0.24D \quad 0.79D \quad 0.95D \quad 1.04D$

It is NCERT Data Based

52. Identify the number of structure/s from the following which can be correlated to D-glyceraldehyde.



(1) three





(2) two

(3) four

(4) one

Ans. (1)

Sol.



In A, B, D – OH group in right hand side then D-configuration is assigned

53. The maximum covalency of a non-metallic group

15 element 'E' with weakest E–E bond is :

(1) 5

(2) 3

(3) 6

(4) 4


Ans. (4)

Sol. N – N < P – P : single (σ) bond strength

Due to L.P.-L.P. repulsion

and maximum possible covalency of nitrogen is 4.

54. Consider the given figure and choose the **correct** option :



- (1) Activation energy of backward reaction is E_1 and product is more stable than reactant.
- (2) Activation energy of forward reaction is $E_1 + E_2$ and product is more stable than reactant.
- (3) Activation energy of forward reaction is $E_1 + E_2$ and product is less stable than reactant.
- (4) Activation energy of both forward and backward reaction is $E_1 + E_2$ and reactant is more stable than product.

Ans. (3)





Sol. Activation energy of forward reaction = $E_1 + E_2$

Energy of product > Energy of reactant


Stability

Reactant > Product

55. When sec-butylcyclohexane reacts with bromine in the presence of sunlight, the major product is :



Ans. (4)



Formation of more stable free radical intermediate

56. The species which does not undergo disproportionation reaction is :

- (1) ClO_2
- (2) ClO_4^-
- (3) ClO^-
- (4) ClO_3^-

Ans. (2)

Sol. $\text{ClO}_4^- \rightarrow x + \{(-2) \times 4\} = -1 \Rightarrow x = +7$

Chlorine is in its maximum oxidation state, so disproportionation not possible in ClO_4^- .


57. Match the Compounds (**List-I**) with the appropriate Catalyst/Reagents (**List-II**) for their reduction into corresponding amines.

	List-I (Compounds)	List-II (Catalyst/Reagents)
(A)	(I)	NaOH (aqueous)
(B)	(II)	H_2/Ni
(C)	(III)	$\text{LiAlH}_4, \text{H}_2\text{O}$
(D)	(IV)	Sn, HCl

Choose the **correct** answer from the options given below :


- (1) (A)-(III), (B)-(II), (C)-(IV), (D)-(I)
- (2) (A)-(II), (B)-(IV), (C)-(III), (D)-(I)
- (3) (A)-(II), (B)-(I), (C)-(III), (D)-(IV)
- (4) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)

Ans. (4)



The maximum number of RBr producing 2-methylbutane by above sequence of reactions is . (Consider the structural isomers only)

Ans. (1)



Sol.  Total number of possible structures of $RBr = 4$

59. Match List-I with List-II.

	List-I (Partial Derivatives)		List-II (Thermodynamic Quantity)
(A)	$\left(\frac{\partial G}{\partial T} \right)_P$	(I)	C _P
(B)	$\left(\frac{\partial H}{\partial T} \right)_P$	(II)	-S
(C)	$\left(\frac{\partial G}{\partial P} \right)_T$	(III)	C _V
(D)	$\left(\frac{\partial U}{\partial T} \right)_V$	(IV)	V

Choose the **correct** answer from the options given below :

- (1) (A)-(II), (B)-(I), (C)-(III), (D)-(IV)
- (2) (A)-(II), (B)-(I), (C)-(IV), (D)-(III)
- (3) (A)-(I), (B)-(II), (C)-(IV), (D)-(III)
- (4) (A)-(II), (B)-(III), (C)-(I), (D)-(IV)

Ans. (2)

Sol. (A) $dG = VdP - SdT$

Constant pressure

$$\left(\frac{\partial G}{\partial T} \right)_p = -S$$

$$(B) dH = (dq)_P = nC_P dT$$

$$\left(\frac{\partial H}{\partial T} \right)_p = C_p$$

$$(C) dG = VdP - SdT$$

At constant temperature

$$dG = V dP$$

$$(D) dU \equiv nC_V dT \equiv (q)_V$$


$$\left(\frac{\partial U}{\partial T} \right)_V = C_V$$

60. The correct order of the following complexes in terms of their crystal field stabilization energies is :

- (1) $[\text{Co}(\text{NH}_3)_4]^{2+} < [\text{Co}(\text{NH}_3)_6]^{2+} < [\text{Co}(\text{en})_3]^{3+} < [\text{Co}(\text{NH}_3)_6]^{3+}$
- (2) $[\text{Co}(\text{NH}_3)_4]^{2+} < [\text{Co}(\text{NH}_3)_6]^{2+} < [\text{Co}(\text{NH}_3)_6]^{3+} < [\text{Co}(\text{en})_3]^{3+}$
- (3) $[\text{Co}(\text{NH}_3)_6]^{2+} < [\text{Co}(\text{NH}_3)_6]^{3+} < [\text{Co}(\text{NH}_3)_4]^{2+} < [\text{Co}(\text{en})_3]^{3+}$
- (4) $[\text{Co}(\text{en})_3]^{3+} < [\text{Co}(\text{NH}_3)_6]^{3+} < [\text{Co}(\text{NH}_3)_6]^{2+} < [\text{Co}(\text{NH}_3)_4]^{2+}$

Ans. (2)

Sol. Order of CFSE



SFL : NH₃ < en

66. Given below are two statements

Statement (I) : Nitrogen, sulphur, halogen and phosphorus present in an organic compound are detected by Lassaigne's Test.

Statement (II) : The elements present in the compound are converted from covalent form into ionic form by fusing the compound with Magnesium in Lassaigne's test.

In the light of the above statements, choose the **correct** answer from the options given below :

- Both Statement I and Statement II are true
- Both Statement I and Statement II are false
- Statement I is true but Statement II is false
- Statement I is false but Statement II is true

Ans. (3)

Sol. The elements present in the compound are converted from covalent form into ionic form by fusing the compound with sodium in Lassaigne's test

67. Identify the homoleptic complex(es) that is/are low spin.

- $[\text{Fe}(\text{CN})_5\text{NO}]^{2-}$
- $[\text{CoF}_6]^{3-}$
- $[\text{Fe}(\text{CN})_6]^{4-}$
- $[\text{Co}(\text{NH}_3)_6]^{3+}$
- $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$

Choose the **correct** answer from the options given below :

- (B) and (E) only
- (A) and (C) only
- (C) and (D) only
- (C) only

Ans. (3)



Sol. (A) $[\text{Fe}(\text{CN})_5\text{NO}]^{2-} \rightarrow$ Heteroleptic, Fe^{+2} , $3d^6$, $t_{2g}^6 e_g^0, d^2sp^3$, Low spin (3d series + SFL)

(B) $[\text{CoF}_6]^{3-} \rightarrow$ Homoleptic, sp^3d^2 , High spin, Co^{+3} , $3d^6$ (3d series + WFL)

(C) $[\text{Fe}(\text{CN})_6]^{4-} \rightarrow$ Homoleptic
 Fe^{+2} , $3d^6$, d^2sp^3 , $t_{2g}^6 e_g^0$ Low spin
(3d series + SFL)


(D) $[\text{Co}(\text{NH}_3)_6]^{3+} \rightarrow$ Homoleptic, Co^{+3} $3d^6$, d^2sp^3 , $t_{2g}^6 e_g^0$, Low spin (3d series + SFL)

(E) $[\text{Cr}(\text{H}_2\text{O})_6]^{2+} \rightarrow$ Homoleptic
 Cr^{+2} $3d^4$, d^2sp^3 , High spin $t_{2g}^3 e_g^1$
(3d series + WFL)




Structure of residue (A) and compound (B)

Formed respectively is :



Ans. (4)



69. Given below are two statements :

Statement (I) : Corrosion is an electrochemical phenomenon in which pure metal acts as an anode and impure metal as a cathode.

Statement (II) : The rate of corrosion is more in alkaline medium than in acidic medium.

In the light of the above statements, choose the **correct** answer from the options given below :

- Both Statement I and Statement II are false
- Statement I is false but Statement II is true
- Both Statement I and Statement II are true
- Statement I is true but Statement II is false

Ans. (4)

Sol. Statement I :

Corrosion is an example of electrochemical phenomenon

In which pure metal act as anode and impure metal (rusted metal) act as cathode.

Statement II :

Corrosion is more favourable in acid medium than alkaline so rate of corrosion is high in acid medium than alkaline.

70. The alkane from below having two secondary hydrogens is :


- (1) 4-Ethyl-3,4-dimethyloctane
- (2) 2,2,4,4-Tetramethylhexane
- (3) 2,2,3,3-Tetramethylpentane
- (4) 2,2,4,5-Tetramethylheptane

Ans. (3)

Alkane	2°H
1.	10
2.	4
3.	2
4.	4

SECTION-B

71. The compound with molecular formula C_6H_6 , which gives only one monobromo derivative and takes up four moles of hydrogen per mole for complete hydrogenation has _____ π electrons.

Ans. (8)

72. Niobium (Nb) and ruthenium (Ru) have "x" and "y" number of electrons in their respective 4d orbitals. The value of $x + y$ is _____

Ans. (11)

$$\text{Sol. } Z = 41 \rightarrow \text{Nb (Niobium)} : [\text{Kr}]_{36} 4d^4 5s^1$$

Number of electron in 4d = 4 = x

$$Z = 44 \rightarrow \text{Ru (Ruthenium)} [\text{Kr}]_{36} 4d^7 5s^1$$


Number of electron in 4d = 7 = y

$$x + y = 11$$

73. The complex of Ni^{2+} ion and dimethyl glyoxime contains _____ number of Hydrogen (H) atoms.




Ans. (14)

$$\text{Sol. } [\text{Ni}(\text{dmg})_2]$$





Number of H-atom = 14


74. Consider the following cases of standard enthalpy of reaction (ΔH_f° in kJ mol^{-1})



The magnitude of $\Delta H_f^\circ \text{C}_2\text{H}_6(\text{g})$ is _____ kJ mol^{-1} (Nearest integer).

Ans. (95)

$$\Delta H_1 = -1550$$



$$\Delta H_f = 2\Delta H_2 + 3\Delta H_3 - \Delta H_1 = 95 \text{ kJ/mole.}$$

75. 20 mL of 2 M NaOH solution is added to 400 mL of 0.5 M NaOH solution. The final concentration of the solution is _____ $\times 10^{-2}$ M. (Nearest integer).

Ans. (57)

$$\text{Sol. } M_F = \frac{M_1 V_1 + M_2 V_2}{V_1 + V_2}$$

$$= \frac{2 \times 20 + 0.5 \times 400}{420} = 0.571 \text{ M}$$

$$= 57.1 \times 10^{-2} \text{ M}$$

$$= 57$$